

YTUMB 00-01, May 2001

Differential Geometry of the q -Quaternions

Salih Çelik¹

Yildiz Technical University, Department of Mathematics,
34210 Davutpasa, Istanbul, TURKEY.

Abstract

Differential calculus on the quantum quaternionic group $GL(1, H_q)$ is introduced.

¹E-mail: sacelik@yildiz.edu.tr

1. Introduction

Differential geometry in the theory of (quantum) Lie groups plays an important role in the mathematical modelling of physics theories. In the classical differential geometry one has a choice between two dual and equivalent descriptions: one can either work with points on a manifold M or with the algebra $C(M)$ of smooth functions on M . The idea that the algebra $C(M)$ need not be commutative gives rise to the noncommutative geometry. Such a space is called a quantum space. This is analogy with the quantization of the commutative algebra of functions on phase space that yields the noncommutative operator algebra of quantum mechanics.

A class of noncommutative Hopf algebras have been found in the discussions of integrable systems. These Hopf algebras are q -deformed function algebras of classical groups and this structure is called quantum group [1]. The quantum group can also be regarded as a generalization of the notion of a group [2]. Noncommutative geometry [3] is one of the most attractive mathematical concepts in physics and has started to play an important role in different fields of mathematical physics for the last few years. The basic structure giving a direction to the noncommutative geometry is a differential calculus [4] on an associative algebra.

Quantum quaternionic algebra and its Hopf algebra structure is important in physics. The importance of differential geometry in the quantum quaternionic algebra should not be underestimated. Quantum quaternion is an example of a quantum space, and to investigate its differential geometry may be interesting. This is considered in the present work.

2. Review of Hopf algebra H_q

Elementary properties of quantum quaternionic group $\text{GL}(1, H_q) = H_q$ are described in Refs. 5 and 6. We state briefly the properties we are going to need in this work.

2.1 The algebra of functions on H_q

The quantum quaternionic algebra H_q is defined as a pair $(\mathcal{A}, \mathcal{M})$ equipped with $*$ structure, where $\mathcal{A}$ is an algebra and $\mathcal{M}$ is an $\mathcal{A}$ -module, and they have the following properties:

(1) $\mathcal{A}$ is an unital associative algebra generated by generators a_k ($k = 0, 1, 2, 3$) with the commutation relations

$$\begin{aligned}
 a_0 a_1 &= a_1 a_0 - \mathbf{i} \frac{q - q^{-1}}{2} (a_2^2 + a_3^2), \\
 a_0 a_2 &= \frac{q + q^{-1}}{2} a_2 a_0 + \mathbf{i} \frac{q - q^{-1}}{2} a_2 a_1, \\
 a_0 a_3 &= \frac{q + q^{-1}}{2} a_3 a_0 + \mathbf{i} \frac{q - q^{-1}}{2} a_3 a_1, \\
 a_1 a_2 &= \frac{q + q^{-1}}{2} a_2 a_1 - \mathbf{i} \frac{q - q^{-1}}{2} a_2 a_0, \\
 a_1 a_3 &= \frac{q + q^{-1}}{2} a_3 a_1 - \mathbf{i} \frac{q - q^{-1}}{2} a_3 a_0, \\
 a_2 a_3 &= a_3 a_2,
 \end{aligned} \tag{1}$$

where $\mathbf{i}^2 = -1$ and q is a nonzero real number.

(2) The $*$ antiinvolution in $\mathcal{A}$ is defined by

$$\begin{aligned}
 a_j^* &= a_j, \quad j = 0, 1 \\
 a_2^* &= \frac{1}{2} \left[(q + q^{-1}) a_2 - \mathbf{i} (q - q^{-1}) a_3 \right], \\
 a_3^* &= \frac{\mathbf{i}}{2} \left[(q - q^{-1}) a_2 - \mathbf{i} (q + q^{-1}) a_3 \right].
 \end{aligned} \tag{2}$$

Note that

$$(a_k^*)^* = a_k, \quad k = 0, 1, 2, 3.$$

(3) $\mathcal{M}$ is an $\mathcal{A}$ -module generated by the quaternionic units e_k with the relations

$$e_k e_l = -\delta_{kl} e_0 + \epsilon_{klm} e_m \quad (3)$$

and

$$a_k e_l = e_l a_k, \quad (4)$$

where δ_{kl} denotes the Kronecker delta and

$$\epsilon_{klm} = \frac{1}{2}(k-l)(l-m)(m-k).$$

The quaternionic conjugation ($\star$ -antiinvolution) in $\mathcal{M}$ is defined by

$$e_k^\star = 2\delta_{0,k} e_0 - e_k, \quad k = 0, 1, 2, 3. \quad (5)$$

(4) Assume that any quaternion h has a representation

$$h = a_0 e_0 + a_1 e_1 + a_2 e_2 + a_3 e_3 \quad (6)$$

in terms of the generators of $\mathcal{A}$. h will be called the q -quaternion, and in this case we shall say that q -quaternion h belongs to H_q .

The conjugation of a q -quaternion h is introduced as

$$h^\star = e_0 a_0^\star - e_1 a_1^\star - e_2 a_2^\star - e_3 a_3^\star. \quad (7)$$

Hence, we can introduce the q -norm of h as

$$\mathcal{N}_q(h) = hh^\star = a_0^2 + a_1^2 + \frac{1}{2}(q + q^{-1})(a_2^2 + a_3^2). \quad (8)$$

Note that $\mathcal{N}_q(h)$ belongs to the center of H_q .

2. 2 The Hopf Algebra Structure of H_q

The action of comultiplication Δ on the generators a_k of $\mathcal{A}$ can be introduced as

$$\begin{aligned} \Delta(a_0) &= a_0 \otimes a_0 - (a_1 \otimes a_1 + a_2 \otimes a_2 + a_3 \otimes a_3), \\ \Delta(a_1) &= a_0 \otimes a_1 + a_1 \otimes a_0 + a_2 \otimes a_3 - a_3 \otimes a_2, \end{aligned} \quad (9)$$

$$\Delta(a_2) = a_0 \otimes a_2 + a_2 \otimes a_0 + a_3 \otimes a_1 - a_1 \otimes a_3,$$

$$\Delta(a_3) = a_0 \otimes a_3 + a_3 \otimes a_0 + a_1 \otimes a_2 - a_2 \otimes a_1.$$

Note that

$$\Delta(e_0) = e_0 \otimes e_0. \quad (10)$$

It also easy to show that

$$\Delta(\mathcal{N}_q(h)) = \mathcal{N}_q(h) \otimes \mathcal{N}_q(h). \quad (11)$$

The action of counit ε on the generators a_k of $\mathcal{A}$ is given by

$$\varepsilon(a_k) = \delta_{0,k}e_0, \quad k = 0, 1, 2, 3 \quad (12)$$

$$\varepsilon(e_0) = e_0.$$

The action of antipode $\mathcal{S}$ on the generators a_k of $\mathcal{A}$ is introduced as

$$\mathcal{S}(a_k) = \mathcal{N}_q^{-1}(h)(2\delta_{0,k}a_0 - a_k^*) \quad (13)$$

for $k = 0, 1, 2, 3$.

3. Differential calculus on H_q

In this section, we shall build up the differential calculus on the quantum quaternionic algebra H_q . The differential calculus on H_q involves functions on the algebra, differentials and differential forms.

3.1 Classical case

Let's begin with differential calculus on the classical quaternionic group $\text{GL}(1, H)$. In a classical Lie group G , one-order differential calculus is a linear map

$$\mathbf{d} : C^\infty(G) \longrightarrow \Gamma,$$

where $C^\infty(G)$ is a $\mathcal{C}$ -algebra consisting of all smooth functions on G and Γ is a $C^\infty(G)$ -bimodule consisting of all differential forms. This linear map satisfies

(i) the nilpotency

$$\mathbf{d}^2 = 0, \quad (14)$$

(ii) for all $f, g \in C^\infty(G)$,

$$\mathbf{d}(fg) = (\mathbf{d}f) g + (-1)^{\hat{f}} f (\mathbf{d}g) \quad (15)$$

where f and g are functions of the generators and $\hat{f}$ is the corresponding grading of f .

According to the ideas of noncommutative geometry [3], the differential calculus can be defined on a more general noncommutative algebra.

Let $\mathcal{A}_1$ be a Hopf algebra with unit generated by the generators a_k ($k = 0, 1, 2, 3$). We denote differentials of a_k by $\mathbf{d}a_k$. Then one can construct the one-form Ω , where

$$\Omega = \mathbf{d}h h^*. \quad (16)$$

Explicitly,

$$w_0 = \mathbf{d}a_0 a_0 + \mathbf{d}a_1 a_1 + \mathbf{d}a_2 a_2 + \mathbf{d}a_3 a_3, \quad (17)$$

etc. It can be easily checked that these one-forms construct a four dimensional Grassmann algebra. The anti-commutation relations of the one-forms allow us to construct the algebra of the generators. To obtain the Lie algebra of the algebra generators we first write the one-forms as

$$\mathbf{d}a_0 = w_0 a_0 - w_1 a_1 - w_2 a_2 - w_3 a_3, \quad (18)$$

etc. The differential $\mathbf{d}$ can then be expressed in the form

$$\mathbf{d} = 2(w_0 \nabla_0 - w_1 \nabla_1 + w_2 \nabla_2 - w_3 \nabla_3), \quad (19)$$

where ∇_k ($k = 0, 1, 2, 3$) are the Lie algebra generators. We wish to obtain the commutation relations of these generators. Let f be an arbitrary function of the generators of $\mathcal{A}_1$. Then, using the nilpotency of the exterior differential $\mathbf{d}$ we obtain

$$(-1)^k \mathbf{d}w_k \nabla_k f = (-1)^{k+j} w_k (w_j \nabla_j) \nabla_k f, \quad k, j = 0, 1, 2, 3 \quad (20)$$

where summation over repeated indices is understood. Using one-forms one easily obtain the two-forms

$$\mathbf{d}w_0 = 0, \quad \mathbf{d}w_1 = 2w_2 w_3,$$

$$\mathrm{d}w_2 = 2w_3w_1, \quad \mathrm{d}w_3 = -2w_2w_1, \quad (21)$$

since

$$\mathrm{d}\Omega = -\mathrm{d}h \, \mathrm{d}h^* = \Omega^2. \quad (22)$$

We now find the following commutation relations for the Lie algebra

$$\begin{aligned} [\nabla_1, \nabla_0] &= [\nabla_2, \nabla_0] = [\nabla_3, \nabla_0] = 0, \\ [\nabla_1, \nabla_2] &= -2\nabla_3, \quad [\nabla_2, \nabla_3] = -2\nabla_1, \quad [\nabla_3, \nabla_1] = -2\nabla_2. \end{aligned} \quad (23)$$

3.2 Quantum case

A differential algebra on H_q is an associative algebra Γ equipped with an operator d . Also the algebra Γ has to be generated by $\mathcal{A} \cup \mathrm{d}\mathcal{A}$.

Firstly, to obtain the relations between the generators of $\mathcal{A}$ and their differentials. We shall use the method of Ref. 7. Using the consistency of a differential calculus, as the final result one has the following commutation relations

$$\begin{aligned} a_0 \, \mathrm{d}x_+ &= \frac{q^2+1}{2} \mathrm{d}x_+ a_0 + \mathbf{i} \frac{q^2-1}{2} \mathrm{d}x_+ a_1, \\ a_1 \, \mathrm{d}x_+ &= \frac{q^2+1}{2} \mathrm{d}x_+ a_1 - \mathbf{i} \frac{q^2-1}{2} \mathrm{d}x_+ a_0, \\ a_0 \, \mathrm{d}x_- &= \frac{q^2+1}{2} \mathrm{d}x_- a_0 - \mathbf{i} \frac{q^2-1}{2} \mathrm{d}x_- a_1 + \frac{(q-q^{-1})^2}{2} \mathrm{d}x_+ x_- - (q-q^{-1})(\mathrm{d}a_2 a_2 + \mathrm{d}a_3 a_3), \\ a_1 \, \mathrm{d}x_- &= \frac{q^2+1}{2} \mathrm{d}x_- a_1 + \mathbf{i} \frac{q^2-1}{2} \mathrm{d}x_- a_1 - \mathbf{i} \frac{(q-q^{-1})^2}{2} \mathrm{d}x_+ x_- + \mathbf{i}(q-q^{-1})(\mathrm{d}a_2 a_2 + \mathrm{d}a_3 a_3), \\ a_0 \, \mathrm{d}a_2 &= q \mathrm{d}a_2 a_0 + \frac{q^2-1}{2} (\mathrm{d}a_0 + \mathbf{i} \mathrm{d}a_1) a_2, \\ a_0 \, \mathrm{d}a_3 &= q \mathrm{d}a_3 a_0 + \frac{q^2-1}{2} (\mathrm{d}a_0 + \mathbf{i} \mathrm{d}a_1) a_3, \\ a_1 \, \mathrm{d}a_2 &= q \mathrm{d}a_2 a_1 - \mathbf{i} \frac{q^2-1}{2} (\mathrm{d}a_0 + \mathbf{i} \mathrm{d}a_1) a_2, \\ a_1 \, \mathrm{d}a_3 &= q \mathrm{d}a_3 a_1 - \mathbf{i} \frac{q^2-1}{2} (\mathrm{d}a_0 + \mathbf{i} \mathrm{d}a_1) a_3, \\ a_2 \, \mathrm{d}a_0 &= q \mathrm{d}a_0 a_2 + \frac{q^2-1}{2} \mathrm{d}a_2 (a_0 - \mathbf{i} a_1), \\ a_2 \, \mathrm{d}a_1 &= q \mathrm{d}a_1 a_0 - \mathbf{i} \frac{q^2-1}{2} \mathrm{d}a_2 (a_0 - \mathbf{i} a_1), \end{aligned} \quad (24)$$

$$\begin{aligned}
a_2 \, \mathbf{d}a_2 &= \frac{q^2+1}{2} \mathbf{d}a_2 \, a_2 - \frac{q^2-1}{2} \mathbf{d}a_3 \, a_3 - \frac{q-q^{-1}}{2} \mathbf{d}x_+ \, x_-, \\
a_2 \, \mathbf{d}a_3 &= \frac{q^2+1}{2} \mathbf{d}a_3 \, a_2 + \frac{q^2-1}{2} \mathbf{d}a_2 \, a_3, \\
a_3 \, \mathbf{d}a_0 &= q \mathbf{d}a_0 \, a_3 + \frac{q^2-1}{2} \mathbf{d}a_3 (a_0 - \mathbf{i}a_1), \\
a_3 \, \mathbf{d}a_1 &= q \mathbf{d}a_1 \, a_3 - \mathbf{i} \frac{q^2-1}{2} \mathbf{d}a_3 (a_0 - \mathbf{i}a_1), \\
a_3 \, \mathbf{d}a_2 &= \frac{q^2+1}{2} \mathbf{d}a_2 \, a_3 + \frac{q^2-1}{2} \mathbf{d}a_3 \, a_2, \\
a_3 \, \mathbf{d}a_3 &= \frac{q^2+1}{2} \mathbf{d}a_3 \, a_3 - \frac{q^2-1}{2} \mathbf{d}a_2 \, a_2 - \frac{q-q^{-1}}{2} \mathbf{d}x_+ \, x_-,
\end{aligned}$$

where

$$\mathbf{d}x_{\pm} = \mathbf{d}a_0 \pm \mathbf{i} \mathbf{d}a_1.$$

Applying the exterior differential $\mathbf{d}$ on the relations (24) and using the nilpotency of $\mathbf{d}$ one obtains

$$\begin{aligned}
\mathbf{d}a_0 \, \mathbf{d}a_1 &= -\mathbf{d}a_1 \mathbf{d}a_0, \\
(\mathbf{d}a_0)^2 &= 0 = (\mathbf{d}a_1)^2, \\
\mathbf{d}a_0 \, \mathbf{d}a_2 &= -\frac{q+q^{-1}}{2} \mathbf{d}a_2 \, \mathbf{d}a_0 + \mathbf{i} \frac{q-q^{-1}}{2} \mathbf{d}a_2 \, \mathbf{d}a_1, \\
\mathbf{d}a_0 \, \mathbf{d}a_3 &= -\frac{q+q^{-1}}{2} \mathbf{d}a_3 \, \mathbf{d}a_0 + \mathbf{i} \frac{q-q^{-1}}{2} \mathbf{d}a_3 \, \mathbf{d}a_0, \\
\mathbf{d}a_1 \, \mathbf{d}a_2 &= -\frac{q+q^{-1}}{2} \mathbf{d}a_2 \, \mathbf{d}a_1 - \mathbf{i} \frac{q-q^{-1}}{2} \mathbf{d}a_2 \, \mathbf{d}a_0, \\
\mathbf{d}a_1 \, \mathbf{d}a_3 &= -\frac{q+q^{-1}}{2} \mathbf{d}a_3 \, \mathbf{d}a_1 - \mathbf{i} \frac{q-q^{-1}}{2} \mathbf{d}a_3 \, \mathbf{d}a_1, \\
\mathbf{d}a_2 \, \mathbf{d}a_3 &= -\mathbf{d}a_3 \, \mathbf{d}a_2, \\
(\mathbf{d}a_2)^2 &= \mathbf{i}(q-q^{-1}) \mathbf{d}a_1 \, \mathbf{d}a_0 = (\mathbf{d}a_3)^2.
\end{aligned} \tag{25}$$

There is an interesting case which gives rise to the second kind of quaternionic variables, Grassmann quaternion is defined by

$$\psi = \psi_0 e_0 + \psi_1 e_1 + \psi_2 e_2 + \psi_3 e_3,$$

where components ψ_k ($k = 0, 1, 2, 3$) are Grassmann variables. Essentially the relations (25) are the relations between the components ψ_k , in q -deformation. More details will be given in Appendix.

To complete the differential calculus, we need the Cartan-Maurer one-forms. In analogy with the one-forms on a Lie group in classical differential geometry, one can construct the one-form Ω where

$$\Omega = dh \, h^* = w_0 e_0 + w_1 e_1 + w_2 e_2 + w_3 e_3.$$

So we can write the one-forms as follows

$$\begin{aligned} w_0 &= \alpha_0 a_0 + \alpha_1 a_1 + \frac{q + q^{-1}}{2}(\alpha_2 a_2 + \alpha_3 a_3) + \mathbf{i} \frac{q - q^{-1}}{2}(\alpha_3 a_2 - \alpha_2 a_3), \\ w_1 &= -\alpha_0 a_1 + \alpha_1 a_0 - \frac{q + q^{-1}}{2}(\alpha_2 a_3 - \alpha_3 a_2) - \mathbf{i} \frac{q - q^{-1}}{2}(\alpha_2 a_2 + \alpha_3 a_3), \\ w_2 &= \alpha_2 a_0 - \alpha_3 a_1 + \frac{q + q^{-1}}{2}(\alpha_1 a_3 - \alpha_0 a_2) + \mathbf{i} \frac{q - q^{-1}}{2}(\alpha_0 a_3 + \alpha_1 a_2), \\ w_3 &= \alpha_2 a_1 + \alpha_3 a_0 - \frac{q + q^{-1}}{2}(\alpha_0 a_3 + \alpha_1 a_2) + \mathbf{i} \frac{q - q^{-1}}{2}(\alpha_1 a_3 - \alpha_0 a_2), \end{aligned} \quad (26)$$

where $\alpha_0 = da_0$, etc.

We wish to find the commutation relations of the generators of $\mathcal{A}$ with those of the components of Ω which may be computed directly, as follows:

$$\begin{aligned} a_0 w_+ &= \frac{q^2 + 1}{2} w_+ a_0 + \mathbf{i} \frac{q^2 - 1}{2} w_+ a_1, & a_2 w_+ &= \frac{q^2 + 1}{2} w_+ a_2 + \mathbf{i} \frac{q^2 - 1}{2} w_+ a_3, \\ a_1 w_+ &= \frac{q^2 + 1}{2} w_+ a_1 - \mathbf{i} \frac{q^2 - 1}{2} w_+ a_0, & a_3 w_+ &= \frac{q^2 + 1}{2} w_+ a_3 - \mathbf{i} \frac{q^2 - 1}{2} w_+ a_2, \\ a_0 w_2 &= q w_2 a_0 + \frac{q - q^{-1}}{2} w_+ a_2, & a_2 w_2 &= q w_2 a_2 - \frac{q - q^{-1}}{2} w_+ a_0, \\ a_1 w_2 &= q w_2 a_1 - \frac{q - q^{-1}}{2} w_+ a_3, & a_3 w_2 &= q w_2 a_3 + \frac{q - q^{-1}}{2} w_+ a_1, \\ a_0 w_3 &= q w_3 a_0 + \frac{q - q^{-1}}{2} w_+ a_3, & a_2 w_3 &= q w_3 a_2 - \frac{q - q^{-1}}{2} w_+ a_1, \\ a_1 w_3 &= q w_3 a_1 + \frac{q - q^{-1}}{2} w_+ a_2, & a_3 w_3 &= q w_3 a_3 - \frac{q - q^{-1}}{2} w_+ a_0, \\ a_0 w_- &= \frac{q^2 + 1}{2} w_- a_0 - \mathbf{i} \frac{q^2 - 1}{2} w_- a_1 + \frac{(q - q^{-1})^2}{2} w_+ (a_0 + \mathbf{i} a_1) + (1 - q^2)(w_2 a_2 + w_3 a_3), \end{aligned} \quad (27)$$

$$\begin{aligned}
a_1 w_- &= \frac{q^2 + 1}{2} w_- a_1 + \mathbf{i} \frac{q^2 - 1}{2} w_- a_0 - \mathbf{i} \frac{(q - q^{-1})^2}{2} w_+ (a_0 + \mathbf{i} a_1) - (1 - q^2)(w_2 a_3 - w_3 a_2), \\
a_2 w_- &= \frac{q^2 + 1}{2} w_- a_2 - \mathbf{i} \frac{q^2 - 1}{2} w_- a_3 + \frac{(q - q^{-1})^2}{2} w_+ (a_2 + \mathbf{i} a_3) + (q^2 - 1)(w_2 a_0 + w_3 a_1), \\
a_3 w_- &= \frac{q^2 + 1}{2} w_- a_3 + \mathbf{i} \frac{q^2 - 1}{2} w_- a_2 - \mathbf{i} \frac{(q - q^{-1})^2}{2} w_+ (a_2 + \mathbf{i} a_3) + (1 - q^2)(w_2 a_1 - w_3 a_0),
\end{aligned}$$

where

$$w_{\pm} = w_0 \pm \mathbf{i} w_1.$$

We now obtain the commutation relations of the Cartan-Maurer forms

$$\begin{aligned}
w_0^2 &= \mathbf{i} \frac{(q - q^{-1})^2}{2} w_3 w_2, & w_1^2 &= \mathbf{i} \frac{q^2 - q^{-2}}{2} w_3 w_2, \\
w_0 w_1 &= -w_1 w_0 + (q^{-2} - 1) w_2 w_3, \\
w_0 w_2 &= -w_2 w_0 + \frac{q^{-2} - q^2}{2} w_3 w_1 + \mathbf{i} \frac{(q - q^{-1})^2}{2} w_2 w_1, \\
w_0 w_3 &= -w_3 w_0 + \frac{q^2 - q^{-2}}{2} w_2 w_1 + \mathbf{i} \frac{(q - q^{-1})^2}{2} w_3 w_1, \\
w_1 w_2 &= -\frac{q^2 + q^{-2}}{2} w_2 w_1 + \mathbf{i} \frac{q^{-2} - q^2}{2} w_3 w_1, \\
w_1 w_3 &= -\frac{q^2 + q^{-2}}{2} w_3 w_1 + \mathbf{i} \frac{q^2 - q^{-2}}{2} w_2 w_1, \\
w_2 w_3 &= -w_3 w_2, & w_2^2 &= 0 = w_3^2.
\end{aligned} \tag{28}$$

The conjugation of one-forms is defined by

$$\begin{aligned}
w_0^* &= q^{-2} w_0 + \mathbf{i} (q^{-2} - 1) w_1, & w_1^* &= w_1, \\
w_2^* &= w_2, & w_3^* &= w_3.
\end{aligned} \tag{29}$$

Note that, we have

$$\Omega + \overline{\Omega} = (1 - q^{-2})(w_0 + \mathbf{i} w_1) e_0. \tag{30}$$

To obtain the quantum algebra of the algebra generators we first write the Cartan-Maurer forms as

$$\begin{aligned}
\mathbf{d}a_0 &= w_0 a_0 - (w_1 a_1 + w_2 a_2 + w_3 a_3), \\
\mathbf{d}a_1 &= w_0 a_1 + w_1 a_0 + w_2 a_3 - w_3 a_2,
\end{aligned}$$

$$\mathbf{d}a_2 = w_0a_2 - w_1a_3 + w_2a_0 + w_3a_1, \quad (31)$$

$$\mathbf{d}a_3 = w_0a_3 + w_1a_2 - w_2a_1 + w_3a_0.$$

Note that

$$\mathbf{d}^\star = q^2 \mathbf{d}. \quad (32)$$

Indeed, for example,

$$\begin{aligned} (\mathbf{d}a_0)^\star &= a_0^\star w_0^\star - a_1^\star w_1^\star - a_2^\star w_2^\star - a_3^\star w_3^\star \\ &= q^{-2}a_0(w_0 + \mathbf{i}w_1) - \mathbf{i}(a_0 - \mathbf{i}a_1)w_1 - \frac{1}{2}a_2[(q + q^{-1})w_2 - \mathbf{i}(q - q^{-1})w_3] \\ &\quad + \frac{\mathbf{i}}{2}a_3[(q - q^{-1})w_2 + \mathbf{i}(q + q^{-1})w_3] \\ &= q^2(w_0a_0 - w_1a_1 - w_2a_2 - w_3a_3) \end{aligned}$$

so that

$$\mathbf{d}^\star a_0^\star = q^2(w_0a_0 - w_1a_1 - w_2a_2 - w_3a_3) = q^2 \mathbf{d}a_0 = q^2 \mathbf{d}a_0^\star$$

implies that

$$\mathbf{d}^\star - q^2 \mathbf{d} = 0.$$

Using the nilpotency of the differential $\mathbf{d}$, we can write the two-forms as

$$\begin{aligned} \mathbf{d}w_0 &= \mathbf{i}(q^{-2} - 1)w_2w_3, & \mathbf{d}w_1 &= (q^{-2} + 1)w_2w_3, \\ \mathbf{d}w_2 &= \mathbf{i}(q^{-2} - 1)w_2w_1 + (q^{-2} + 1)w_3w_1, & (33) \\ \mathbf{d}w_3 &= \mathbf{i}(q^{-2} - 1)w_3w_1 - (q^{-2} + 1)w_2w_1. \end{aligned}$$

Using the Cartan-Maurer equations with together (27) we find the following commutation relations for the quantum algebra:

$$\begin{aligned} \nabla_0 \nabla_1 &= \nabla_1 \nabla_0, & \nabla_0 \nabla_2 &= \nabla_2 \nabla_0, & \nabla_0 \nabla_3 &= \nabla_3 \nabla_0, \\ \nabla_1 \nabla_2 &= \frac{q^2 + q^{-2}}{2} \nabla_2 \nabla_1 - (q^{-2} + 1) \nabla_3 + \mathbf{i}(q^{-2} - 1) \nabla_2 - \mathbf{i} \frac{(q - q^{-1})^2}{2} \nabla_0 \nabla_1 \\ &\quad - \frac{q^2 - q^{-2}}{2} \nabla_3 (\nabla_0 + \mathbf{i} \nabla_1), \end{aligned}$$

$$\begin{aligned}\nabla_1 \nabla_3 &= \frac{q^2 + q^{-2}}{2} \nabla_3 \nabla_1 + (q^{-2} + 1) \nabla_2 - \mathbf{i}(q^{-2} - 1) \nabla_3 - \mathbf{i} \frac{(q - q^{-1})^2}{2} \nabla_0 \nabla_1 \\ &\quad + \frac{q^2 - q^{-2}}{2} \nabla_2 (\nabla_0 + \mathbf{i} \nabla_1),\end{aligned}$$

$$\begin{aligned}\nabla_3 \nabla_2 &= \nabla_2 \nabla_3 + (q^{-2} + 1) \nabla_1 - \mathbf{i}(q^{-2} - 1) \nabla_0 + \mathbf{i} \frac{q^2 - q^{-2}}{2} \nabla_0^2 \\ &\quad + \frac{(q - q^{-1})^2}{2} \nabla_1^2 + (1 - q^{-2}) \nabla_0 \nabla_1.\end{aligned}$$

Acknowledgement

This work was supported in part by **T. B. T. A. K.** the Turkish Scientific and Technical Research Council.

Appendix

q -Deformation of Grassmann quaternionic algebra

Quaternionic Lie groups can be defined in terms of matrices with quaternionic elements. To make contact with the usual formulation of Lie groups in terms of matrices with complex matrix elements, it is useful to represent each quaternion in terms of 2x2 complex matrices. Therefore, the quantum deformation of the Grassmann quaternionic algebra may be introduced using the idea of the quantum matrix theory [2,8].

Quantum Grassmann quaternionic algebra $\hat{H}_q$ is defined as a pair $(\hat{\mathcal{A}}, \mathcal{M})$, where $\hat{\mathcal{A}}$ is an algebra and has the following properties:

$$\begin{aligned}\psi_0^2 &= 0 = \psi_1^2, & \psi_0 \psi_1 + \psi_1 \psi_0 &= 0, \\ \psi_0 \psi_2 &= -\frac{q + q^{-1}}{2} \psi_2 \psi_0 + \mathbf{i} \frac{q - q^{-1}}{2} \psi_2 \psi_1, \\ \psi_0 \psi_3 &= -\frac{q + q^{-1}}{2} \psi_3 \psi_0 + \mathbf{i} \frac{q - q^{-1}}{2} \psi_3 \psi_1, \\ \psi_1 \psi_2 &= -\frac{q + q^{-1}}{2} \psi_2 \psi_1 - \mathbf{i} \frac{q - q^{-1}}{2} \psi_2 \psi_0,\end{aligned}$$

$$\begin{aligned}\psi_1\psi_3 &= -\frac{q+q^{-1}}{2}\psi_3\psi_1 - \mathbf{i}\frac{q-q^{-1}}{2}\psi_3\psi_0, \\ \psi_2^2 &= \mathbf{i}\frac{q-q^{-1}}{2}\psi_1\psi_0 = \psi_3^2, \quad \psi_2\psi_3 + \psi_3\psi_2 = 0.\end{aligned}$$

References

- [1] V. G. Drinfeld, Quantum groups, in *Proc. Int. Cong. Math.*, 798 (Berkeley, 1986).
- [2] N. Y. Reshetikhin, L. A. Takhtajan and L. D. Faddeev, Quantisation of Lie groups and Lie algebras, *Leningrad Math. J.* **1** (1990), 193;
Yu I. Manin, Quantum groups and noncommutative geometry, (Montreal Univ. 1988).
- [3] A. Connes, Noncommutative Geometry, (Academic Press, 1991).
- [4] S. L. Woronowicz, Differential calculus on compact matrix pseudogroups, *Commun. Math. Phys.* **122** (1989), 125;
S. L. Woronowicz, Twisted $SU(2)$ group. An example of a noncommutative differential calculus, *Publ. Res. Inst. Math. Sci.* **23** (1987), 117;
J. Wess and B. Zumino, *Nucl. Phys. (Proc. Suppl.)* **18 B** (1990), 302.
- [5] S. Marchiafava and J. Rembielinski, Quantum quaternions, *J. Math. Phys.* **33** (1992), 171.
- [6] S. Çelik, The Hopf algebra structure of $GL(1, H_q)$ and the isomorphisms $SP_q(1)$ and $SU_q(2)$, *Lett. Math. Phys.* **42** (1997), 27;
S. Çelik, Quantum Matrix Groups and q -Oscillators, (in Turkish) Ph. D. Thesis, (Istanbul Technical University, 1992).
- [7] S. Çelik and S. A. Çelik, On the differential geometry of $GL_q(1|1)$, *J. Phys. A* **31** (1998), 9685.
- [8] S. Çelik, Two parameter deformation of grassmann matrix groups and supergroups, *J. Math. Phys.* **37** (1996), 3568.